

NONTRIVIAL RATIONAL POLYNOMIALS IN TWO VARIABLES HAVE REDUCIBLE FIBRES

WALTER D. NEUMANN AND PAUL NORBURY

We shall call a polynomial map $f: \mathbb{C}^2 \rightarrow \mathbb{C}$ a “coordinate” if there is a g such that $(f, g): \mathbb{C}^2 \rightarrow \mathbb{C}^2$ is a polynomial automorphism. Equivalently, by Abhyankar-Moh and Suzuki, f has one and therefore all fibres isomorphic to $\mathbb{C}$. Following [7] we call a polynomial $f: \mathbb{C}^2 \rightarrow \mathbb{C}$ “rational” if the general fibres of f (and hence all fibres of f) are rational curves. The following theorem, which says that a rational polynomial map with irreducible fibres cannot be part of a counterexample to the 2-dimensional Jacobian Conjecture, has appeared in the literature several times. It appears with an algebraic proof in Razar [12]. It is Theorem 2.5 of Heitmann [4] (as corrected in the Corrigendum), and Lê and Weber, who give a geometric proof in [6], also cite the reference Friedland [3], which we have not seen.

Theorem 1. *If $f: \mathbb{C}^2 \rightarrow \mathbb{C}$ is a rational polynomial map with irreducible fibres and is not a coordinate then f has no jacobian partner (i.e., no polynomial g such that the jacobian of (f, g) is a non-zero constant).*

In this note we prove the above theorem is empty:

Theorem 2. *There is no f satisfying the assumptions of the above theorem. That is, a rational f with irreducible fibres is a coordinate.*

Proof. This theorem is implicit in [7]. Suppose f is rational. As in [7], [6], [9], etc., we consider a nonsingular compactification $Y = \mathbb{C}^2 \cup E$ of $\mathbb{C}^2$ such that f extends to a holomorphic map $\bar{f}: Y \rightarrow \mathbb{P}^1$. Then E is a union of smooth rational curves $E_1, \dots, E_n$ with normal crossings. An E_i is called *horizontal* if $\bar{f}|_{E_i}$ is non-constant. Let δ be the number of horizontal curves. Then we have

$$\delta - 1 = \sum_{a \in \mathbb{C}} (r_a - 1),$$

where r_a is the number of irreducible components of $f^{-1}(a)$. This is Lemma 1.6 of Miyanishi and Sugie [7] who attribute it to Saito [10] and Lemma 4 of Lê-Weber [6] who attribute it to Kaliman [5], corollary 2. The proof is simple arithmetic from the topological observation that on the one hand the euler characteristic of Y is $n + 2$ and on the other hand it is $4 + \sum_{a \in \mathbb{P}^1} (\bar{r}_a - 1)$, where $\bar{r}_a$ is the number of components of $\bar{f}^{-1}(a)$, $a \in \mathbb{P}^1$.

By this formula, if f has irreducible fibres then there is just one horizontal curve. Lemma 1.7 of [7] now says that f is a coordinate. This also follows from the following proposition, which implies that the generic fibres of f have just one point at infinity and are thus isomorphic to $\mathbb{C}$. $\square$

This research is supported by the Australian Research Council.

Proposition 3. *Let $f: \mathbb{C}^2 \rightarrow \mathbb{C}$ be any polynomial map and $\bar{f}: Y \rightarrow \mathbb{P}^1$ an extension as above. Denote by d the greatest common divisor of the degrees of $\bar{f}$ on the horizontal curves of Y and D the sum of these degrees. Then the general fibre of f has d components (so $f = h \circ f_1$ for some polynomials $f_1: \mathbb{C}^2 \rightarrow \mathbb{C}$ and $h: \mathbb{C} \rightarrow \mathbb{C}$ with $\deg(h) = d$), each of which is a compact curve with D/d punctures.*

Proof. Let $E_1, \dots, E_\delta$ be the horizontal curves and $d_1, \dots, d_\delta$ be the degrees of $\bar{f}$ on these. Note that the points at infinity of a general fibre $f^{-1}(a)$ are the points where $\bar{f}^{-1}(a)$ meet the horizontal curves E_i , so there are d_i such points on E_i for $i = 1, \dots, \delta$. The relationship between plumbing diagram and splice diagram (cf. [9, 2] says that the splice diagram Γ for a regular link at infinity for f (cf. [8]) has δ nodes with arrows at them, and the number of arrows at these nodes are $d_1, \dots, d_\delta$ respectively. Let Γ_0 be the same splice diagram but with $d_1/d, \dots, d_\delta/d$ arrows at these nodes. Then a minimal Seifert surface S for the link represented by Γ will consist of d parallel copies of a minimal Seifert surface for the link represented by Γ_0 , so this S has d components. But the general fibre of f is such a minimal Seifert surface ([8], Theorem 1), completing the proof. (It also follows that Γ_0 is the regular splice diagram for the polynomial f_1 of the proposition.) $\square$

REFERENCES

- [1] S. Abhyankar and T.T. Moh, Embeddings of the line in the plane, J. Reine Angew. Math. **276** (1975), 148–166.
- [2] D. Eisenbud and W.D. Neumann, “Three dimensional link theory and invariants of plane curve singularities” Ann. Math. Study 101 (Princeton University Press 1985).
- [3] S. Friedland, On the plane jacobian conjecture, Preprint IHES, May 1994 (*per* [6]).
- [4] R. Heitmann, On the Jacobian conjecture, J. Pure Appl. Algebra **64** (1990), 36–72 and Corrigendum *ibid.* **90** (1993), 199–200.
- [5] S. Kaliman, Two remarks on polynomials in two variables, Pacific J. Math. **154** (1992), 285–295.
- [6] Lê Dung Tráng and Claude Weber, Polynômes à fibres rationnelles et conjecture de jacobienne à 2 variables, C. R. Acad. Sci. Paris, **320** (1995), 581–584.
- [7] M. Miyanishi and T. Sugie, Generically rational polynomials, Osaka J. Math. **17** (1980), 339–362.
- [8] W.D. Neumann, Complex algebraic plane curves via their links at infinity, Inv. Math. **98** (1989), 445–489.
- [9] W.D. Neumann, Irregular links at infinity of complex affine plane curves, Quarterly J. Math. (to appear).
- [10] H. Saito, Fonctions entières qui se reduisent à certains polynômes. II, Osaka J. Math. **9** (1977), 649–674.
- [11] M. Suzuki, Propriétés topologiques des polynômes de deux variables complexes, et automorphismes algébriques de l’espace $\mathbb{C}^2$, J. Math. Soc. Japan **26** (1974), 241–257.
- [12] Michael Razar, Polynomial maps with constant Jacobian. Israel J. Math. **32** (1979), 97–106.

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF MELBOURNE, PARKVILLE, VIC 3052, AUSTRALIA

E-mail address: `neumann@maths.mu.oz.au`

DEPARTMENT OF MATHEMATICS, THE UNIVERSITY OF MELBOURNE, PARKVILLE, VIC 3052, AUSTRALIA

E-mail address: `norbs@maths.mu.oz.au`